



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
**Govind Ballabh Pant University of Agriculture & Technology**  
Udham Singh Nagar, Pantnagar, Uttarakhand



## Agromet Advisory Bulletin

Date : 02-12-2022

Weather Forecast of District NAINITAL(Uttarakhand) Issued On : 2022-12-02(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-12-03	2022-12-04	2022-12-05	2022-12-06	2022-12-07
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	18.0	17.0	17.0	17.0	17.0
Tmin(°C)	8.0	7.0	7.0	7.0	7.0
RH-I(%)	75	75	70	70	70
RH-II(%)	35	30	30	30	30
Wind Speed(kmph)	6.0	6.0	6.0	6.0	6.0
Wind Direction(Degree)	230	270	310	320	320
Cloud Cover(Octa)	3	0	0	0	0

### Weather Summary/Alert:

In the coming five days, weather will remain clear. Maximum & minimum temperature may range from 17.0-18.0 deg C & 7.0-8.0 deg C respectively. Wind will blow with a speed of 6.0 km/hr from the SW & NW direction.

### General Advisory:

IMD Weather forecast and Agrometeorological advisory of your location is now available on Meghdoot mobile app in English and regional languages. Kindly download from following link Android:  
<https://play.google.com/store/apps/details?id=com.aas.meghdoot> iOS:  
<https://apps.apple.com/in/app/meghdoot/id1474048155>

### SMS Advisory:

आगामी पांच दिनों में, मौसम साफ रहेगा। किसान भाई रबी फसलों की बुवाई करें व फसलों की बुवाई से पूर्व बीज उपचार अवश्य करें।

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
WHEAT	Sowing of late sown varieties of wheat should be done. Seed should be treated before sowing of rabi crops. Before sowing the wheat, seed treatment should be done with carboxyl 2.5 gram or Tebeconazol 1 g or carbendazim ( bavastin or 50 W P) 2.5 g per kg of seed.
BARLEY	Weeding and hoeing should be done in barley crop.

<b>Crop(Varieties)</b>	<b>Crop Specific Advisory</b>
MUSTARD	In valley and low hills, light irrigation should be given at the time of grain filling.

**Horticulture Specific Advisory:**

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
ONION	In irrigated valley, field preparation should be done for transplanting the seedlings of matured onion and dung manure should be used in good quantity in the field.
POTATO	In irrigated valley, where early varieties of potato have been sown 2-2.5 months ago, digging of potato should be done.
CAULIFLOWER	In irrigated valley, top dressing of urea should be done at the time of weeding and hoeing in transplanted snowball cauliflower.

**Live Stock Specific Advisory:**

<b>Live Stock(Varieties)</b>	<b>Live Stock Specific Advisory</b>
BUFFALO	To save the animal from cold, dry grasses, paddy residue (Puwal) etc which are not used as fodder for animals should be used as a bedding material for animals in sheds. Door and window should be covered properly so that cold wind could not be entered in the animal shed. Sitting place of animals should be leveled. It is advised that green fodder can be given to the animals by mixing it in dry fodder otherwise animal may be infected by Timpeti disease which causes death of animals.