

# Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Govind Ballabh Pant University of Agriculture & Technology

Udham Singh Nagar, Pantnagar, Uttarakhand



#### **Agromet Advisory Bulletin**

Date: 27.09.2024

Weather Forecast of District NAINITAL (Uttarakhand) Issued On: 2024-09-27 (Valid Till 08:30 IST of the next 5 days)

Parameter	28/09/2024	29/09/2024	30/09/2024	01/10/2024	02/10/2024
Rainfall (mm)	30	20	12	8	6
Max Temp (°C)	21	20	21	22	23
Min Temp (°C)	16	15	16	17	17
Max RH I (%)	85	85	75	70	70
Min RH II (%)	40	40	35	30	30
Wind speed (km/h)	4	3	4	4	3
Wind direction	110	50	340	320	320
Cloud cover	7	8	6	4	3

#### **Weather Summary/ Alert:**

The upcoming week predicts light to moderate rainfall of 6-30 mm from 28 September-01 October and maximum-minimum temperature varying between 20.0-23.0oC and 15.0-17.0oC, respectively. The wind is expected to blow from east-south-east, north-east, north-north-west and north-west at a speed of 3-4 kmph. Light to moderate rain likely to occur at a many places on 27 th Sep, 2024, at few places on 28 September and at isolated places from 29 September to 3 October. A yellow warning regarding occurrence of heavy rainfall/thunderstorm accompanied with lightening/ intense to very intense rain spell is expected to occur at isolated places on 27 September and similar conditions without rainfall may continue at isolated places on 28 September.

#### **General Advisory:**

For regular updates on weather conditions in the region, the farmers can get updates from the app "Meghdoot" and lightning update from the app "Damini" available on Google Play Store (Android users) and App Centre (iOS users). The NDVI shows good agricultural vigour in the isolated regions of the state ranging 0.20-0.40. The extended range forecast shows large excess rainfall and normal maximum-minimum temperature trend during 27.09.2024 to 03.10.2024.

#### **SMS Advisory:**

As per IMD forecast light to medium and yellow warning regarding heavy rainfall has been given so farming activities should be scheduled accordingly and proper drainage should be provided.

### **Crop Specific Advisory:**

Crop (Varieties)	Stage	Crop Specific Advisory
Rice	Reproductive stage/	Proper drainage should be provided and farming operations should be done
	Maturity	under clear weather conditions.
Maize	Vegetative/ Maturity	Proper drainage should be provided and farming operations should be done
		under clear weather conditions. The early maize varieties should be
		harvested and harvested produce should be stored well.
Finger millet	Reproductive stage/	Proper drainage should be provided and farming operations should be done
	Maturity	under clear weather conditions
		The matured millet varieties should be harvested and stored well.
Pigeon Pea (Red	Vegetative	Proper drainage should be provided and farming operations should be done
Gram/Arhar)		under clear weather conditions.

	Soyabean	Flowering/pod	Proper drainage should be provided and farming operations should be done
ı		formation	under clear weather conditions.
ı	Green gram	Maturity/Harvesting	Proper drainage should be provided and farming operations should be done
	Black gram		under clear weather conditions. The mature crop should be harvested and
			stored for procurement/ consumption.

## **Horticulture Specific Advisory:**

Horticulture	Stage	Horticulture Specific Advisory
(Varieties)		
Cauliflower	Seedling growth	As per the soil conditions hoeing and weeding should be
		done in the field. Proper drainage should be provided and
		farming operations should be done under clear weather conditions.
Radish/Carrot/ Beetroot	Sowing/Germination	
		be done under clear weather conditions.
Sem/Bakla	Sowing	Proper drainage should be provided and farming operations should
		be done under clear weather conditions.
Citrus	Fruiting	If symptoms of citrus yellow mosaic virus is observed then
		prune the infected twigs and spray systemic insecticide
		such as imidacloprid 17.8 SL 1 ml/3 litre water or
		Thiamethoxam 25% WG @ 1g/3 litre water. Apply first
		spray of Thiamethoxam during initial appearance of pest
		and repeat $2-3$ sprays at $15-21$ days interval depending
		on the level of pest intensity.
		Proper drainage should be provided and farming operations should
		be done under clear weather conditions.

**Live Stock Specific Advisory** 

T !41-	T ! C4- 1- C !!! - A 1-!		
Livestock	Live Stock Specific Advisory		
Cow	After 15 days calves should be fed with calcium, magnesium, phosphorus, zinc, iron, iodine		
	and copper elements should be consumed in the form of salt mixture or mineral drops.		
Goat/Sheep	In rural areas, 2 shots of tetanus toxide should be administered to sheep and goats at one		
	month and the other at 5 months, so that newborn lambs do not get tetanus disease.		

Poultry Specific Advisory

Poultry	Poultry Specific Advisory
Hens	Due to moisture, fungus grows in the feed due to which there is a deficiency of nutrients in the feed, which leads to a disease called aflatoxicosis. Due to this, worms develop in the
	stomach of the hens and their egg-laying capacity decreases. Such hens must be given deworming medicine once a month as per the advice of the veterinarian.